

INTERNATIONAL RESEARCH FELLOWS ASSOCIATION'S


RESEARCH JOURNEY

International E-Research Journal

PEER REFREED & INDEXED JOURNAL

December 2020 Special Issue 256 (C)

Multidisciplinary Research



Guest Editor -

Prof. Dr. Rajani Shikhare,
Principal,
R. B. Attal College, Georai
Dist. - Beed.

Executive Editors :

Dr. B. D. Rupnar,
Dr. P. P. Pangrikar
Mr. S.S. Nagare
Mr. Ranjeet Pagore,

Chief Editor : **Dr. Dhanraj T. Dhangar**

This Journal is indexed in :

- Scientific Journal Impact Factor (SJIF)
- Cosmoc Impact Factor (CIF)
- Global Impact Factor (GIF)
- International Impact Factor Services (IIFS)



'RESEARCH JOURNEY' International E- Research Journal

Impact Factor - (SJIF) - 6.625 (2019),

Special Issue -256 (C) : Multidisciplinary Research

Peer Reviewed Journal

E-ISSN :

2348-7143

Dec. 2020

Impact Factor – 6.625

E-ISSN – 2348-7143

INTERNATIONAL RESEARCH FELLOWS ASSOCIATION'S

RESEARCH JOURNEY

International E-Research Journal

PEER REFREED & INDEXED JOURNAL

December 2020 Special Issue 256 (C)

Multidisciplinary Research

Guest Editor -

Prof. Dr. Rajani Shikhare,
Principal,
R. B. Attal College, Georai
Dist. - Beed.

Executive Editors :

Dr. B. D. Rupnar,
Dr. P. P. Pangrikar
Mr. S.S. Nagare
Mr. Ranjeet Pagore,

Chief Editor : **Dr. Dhanraj T. Dhangar**

Our Editors have reviewed papers with experts' committee, and they have checked the papers on their level best to stop furtive literature. Except it, the respective authors of the papers are responsible for originality of the papers and intensive thoughts in the papers. Nobody can republish these papers without pre-permission of the publisher.

- Chief & Executive Editor

SWATIDHAN INTERNATIONAL PUBLICATIONS

For Details Visit To : www.researchjourney.net

*Cover Photo (Source) : Internet

© All rights reserved with the authors & publisher

Price : Rs. 1000/-



INDEX

No.	Title of the Paper	Author's Name	Page No.
1	Releastic Approach in R. K. Narayan's Novel 'The Guide'	Dr.V. S. Bandal	04
2	Cultural Studies : An Introduction	Mr. Arun Jadhav	11
3	Sanitation and Social Change	Mr. R. B. Kale	13
4	Rotating Fluid of Magneto Hydrodynamics Flow Past an Impulsively Started Infinite Vertical Plate	Vinod Kulkarni, Vijay Sangale	16
5	An Efficient Synthesis of 5-Substituted 1H-Tetrazole Using Eton's Reagent in Water	Rupnar B.D, Shirsat A.J. Jadhav S. B. Bhagat S.S.	22
6	Crop Insurance in India	B.S.Jogdand	27
7	Outline of Modern Research	Dr. Laxmikant Jirewad	31
8	Second ARCs Views on Right to Information Act	Hanmant Helambe	35
9	An Introduction to Smart Libraries	R.B. Pagore, Dr. B. V. Chalukya	38
10	Impact of Cassine Albens Gum on Incidence of Seed Mycoflora in Different Crop Seeds	K.V. Badar, P.P. Pangrikar	47
11	Synthesis, Characterization and Antimicrobial Analysis of Some New Substituted Pyrazoles From Chromones	Amol Shirsat, Balaji Rupnar, Sunil Bhagat	52
12	Synthesis and Characterization of Ni (II) and Mn (II) Metal Complexes of Novel Schiff's Base Ligand	Vrushali Gavhane, Anjali Rajbhoj, Suresh Gaikwad	57
13	Image Classification Using Fuzzy Logic	Pradeep Gaikwad	61
14	Resistivity of Food Preservative Potassium Meta -Bisulphate Using (TDR) Technique	S. G Badhe, S. N. Helambe, T. A.Prajapati	65
15	Studies on Effects of Gamma Radiation on Iron Oxide in the Energy Range 122-1330 Kev	Pradip S. Dahinde	68
16	Effect of N-Fertilizers on Silage Fermentation	Smita Basole , Sunita Bhosle and Prashant Pangrikar	74
17	Investment Awareness Program (IAP): Need in Uncertain Market Conditions	Dr. Sandip Vanjari	79
18	Impact of Covid19 on Health and Hidden Cost of Covid	Dr. Vivek Waykar	83
19	Studies on Physico-Chemical Parameters of Bore Well Water in Satara Parisar, Aurangabad, India	Jagannath Godse, Sanjay Ubale	86
20	Synthesis and Antimicrobial Screening of Novel Pyrazole Substituted Chlorochromones	S. S. Bhagat, B. D. Rupnar, A. J.Shirsat	89
21	Women's Human Rights & Women Empowerment	Dr. S.N. Satale	92
22	Biodiversity of Butterflies Around Georai Region	A. M. Budrukhar	96
23	चूडिया की खनखनाहट और पायलों से फुटते विद्रोह का बिगुल : 'बेघर सपने'	संतोष नागरे	99
24	लोकनाट्य आणि समाजशास्त्र	डॉ. संदीप बनसोडे	105
25	मराठी भाषा आणि साहित्यासाठी एकविसाव्या शतकाची सुरुवात	डॉ. समाधान इंगळे	107
26	दलित स्त्री जीवन के शोषण का जिवंत दस्तावेज : 'जीवन हमारा'	प्रो. रजनी शिखरे, राजाराम जाधव	110



Impact of *Cassine Albens* Gum on Incidence of Seed Mycoflora in Different Crop Seeds

K.V. Badar and P.P. Pangrikar*

Dept of Botany, Yeshwantrao Chavan College, Sillod, Dist. Aurangabad(M.S.) India

Department of Botany, R. B. Attal college, Georai, Dist. Beed*

ABSTRACT :

Gum is naturally occurring chemical substance in the plant. Gums are natural exudates of stems or from the wounds of the plants. Physical ,chemical and Biological properties of gum change as per the eco-climatic condition of the plant.

The gum of Cassine species has been used as incense in religious and cultural ceremonies and in medicines since time immemorial. (Ahmed I. Mehmood Z., Mohammad F. 1998). The cassine albens (R.) belonging to the family Celastraaceae . The common name of Cassine albens is Damebdi. It contains rich chemical substance in the plant Gum. The plant grows in dry mountainous regions of India, Northern Africa and Middle East.. After processing, the gum is then graded according to its flavour, colour, shape and size. In India, the States of Maharashtra ,kerala, Uttar pradesh Madhya Pradesh, Gujarat are the main source of Cassine albens .

There are different name of plants according to locality. Gum is a substance which exudates naturally from the stem or from the wounds of trees. Gums are colloidal in nature soluble in water but completely insoluble in alcohol and ether. Impact of different time period of select plant gum concentration on incidence of seed mycoflora of cereals, pulses and oil seeds like wheat, jowar, cowpea, gram, safflower and soybean. Percent germination or percent inhibition of germination, root and shoot length of seedlings was measured after 7 days of incubation at room temperature. were studied and results shows the impact of Cassine albens gum was found to promote the germination of wheat, jowar, cowpea, and soybean at 10 hour treatment.

Key Words: Gum, *Cassine*, Seed Germination.

Introduction :

The gum-resins of commercial importance collected from the forest are gum karaya, gum ghatti, salai gum, guggul, and gums from various species of Acacia, including Indian gum arabic from *Acacia nilotica* and true gum arabic from *A. senegal*. The importance of commercial resins are obtained from Pinaceae (rosin, amber), Leguminosae (copal) and Dipterocarpaceae (dammar) families.(Babu, R., and Murugan, K. 1998).

The uses of natural gums and resins in food, medicines and in varnishes or as protective coatings go back to very early times. The present day uses of natural gums and resins are numerous and they are employed by a large number of manufacturing industries including food and pharmaceutical industries.

Gums are a group of plant product resembling carbohydrates and widely distributed in vegetable kingdom. The vegetable gum constitute an important group of economic plant product and are utilize in several ways. It is very commonly used by human being from ancient time. Nowadays in modern industry large quantity of gum have been used for manufacture of pharmaceutical products. Gums are characterized by ability to dissolve in water forming viscid solution by



absorbing water to form jelly gelatinous paste. When these jelly exposed to air these gelatinous paste loose the water and dry to hard clear or with different colour Gums are characterized by ability to dissolve in water forming viscid solution by absorbing water to form gelatinous paste. In some cases the production of gum has been attributed to fungi attacking the plant, these fungi being responsible for enzymes that penetrate the tissues and transform the celluloses and hemicelluloses of the cell wall into gum. (Zalucki M.P. and Malcollum S.B. 1999).

Malcolm (1936) concluded that the production of gum in Sudan gum Arabic trees is due to bacterial agency. The real cause of production of gum in many trees is uncertain. The best use of gum is to prepare sticky substance for pasting the paper and other things. Treatment of gums to seeds has positive effect on seed germination. *Acacia* gum is being widely used as an experimental vehicle for drugs in physiological and pharmacological experiments, and it is supposed to be an inert substance, recent reports have confirmed that it has some biological properties as an antioxidant (Trommer and Neubert, 2005; Ali and Al Moundhri, 2006, Hinson *et al.*, 2004) on the metabolism of lipids (Tiss *et al.*, 2001, Evans *et al.*, 1992), positive contribution in treating kidney, (Matsumoto *et al.*, 2006; Bliss *et al.*, 1996, Ali *et al.*, 2008), cardiovascular (Glover *et al.*, 2009) and gastrointestinal diseases (Wapnir *et al.*, 2008, Rehman *et al.*, 2003). *Acacia* gum is usually referred as Gum Arabic (GA). It is an edible biopolymer obtained as exudates of mature trees of *Acacia* spp. The exudate is a non-viscous liquid, rich in soluble fibers, and its emanation from the stems and branches usually occurs under stress conditions such as drought, poor soil fertility, and injury (Williams and Phillips, 1990).

Gums are to be found in a greater or less degree in most plant families. The gum occurs in about 44 families covering 1,900 genera and 21,000 species. Various organs of the plant may produce or secrete them. They may be produced only in very small quantity and not be readily discernible or they may be produced very copiously forming large, conspicuous incrustations on the surface, as with most of the commercial gums, particularly the tree gums. Certain families of flowering plants are notable for the number of species they contain that are free gum yielders. Among them notable is the Leguminosae, in which a hundred or more species of *Acacia* alone are known to yield gum, including those that are commercially important for gum arabic. Several species of *Astragalus* are also free gum yielders and are the source of gum tragacanth. Some additional notable gum yielding genera in the family are *Albiszia*, *Bauhinia*, *Caesalpinia*, *Ceratonia* and *Pithecolobium*. Other important gum yielding families are Anacardiaceae, Combretaceae, Meliaceae, Rosaceae and Rutaceae.

Material And Methods :

01) Collection of Gum samples: Plant gums were regularly collected in all the seasons. It was done by using axe, sterilized blade. Fine cut was made at different parts of the plant, like root, stem, leaves, flower and fruits. Later on at 30, 45 and 60 days exudates gums were collected in presterilized plastic bags, kept in laboratory condition until it was used (Badar, 2011).

02) Preparation of fine powder of Gum: The fine powder of collected dry gums was prepared by using mixer grinder and kept in clean glass pots at 4-6°C temperature.

03) Application of gum on seed germination: The effect of different time period (1 hr, 3 hr, 5hr, 10hr) of plant gum on seed germination and the impact of different concentration like 1%, 2%, 3% and 4% on cereal, pulses and oil seeds of seed mycoflora have been studied. The impact of different time period of percent gum concentration of were studied o *Cassine albensn* the



germination of cereals, pulses and oil seeds like wheat, jowar, cowpea, gram, safflower, soybean and the results are summarized in Table. *Cassine albens* gum was applied on germinating seeds for 1 to 10 hour treatments. The results were recorded at different time intervals by measuring shoot length and root length by vigour formula (Abdul Baki A.A.and Anderson J.D.1975).

Results And Discussion :

It was calculated from the 1% concentration of gum was found *Cassine albens* to promote the germination of wheat, jowar, cowpea and soybean at 10 hour treatment. The gum *Cassine albens* were tested for percent seed germination in table 01 that 1% concentration of gum was found to promote the percent germination of wheat, jowar and cowpea at 1, 3, 5 and 10 hour treatment. The shoot and root length of all treated seed except gram were found to be increased at 1, 3, 5 and 10 hour treatment as compared to control.

1 hour treatment was found to promote the germination of wheat, cowpea, gram and soybean. It was also found to promote the shoot length, root length of wheat, cowpea, gram whereas it was found to retard the growth of shoot and root length of jowar and soybean. Similar type of results were recorded for 3 hour 5 hour treatment. Whereas, at 10 hour treatment the shoot and root length of all treated seeds except Soyabea n were found to be increased as compared to control.

Table 01: Impact of different time period of gum on s *Cassine albens* seed germination (1% gum concentration)

Table with 16 columns: Sr. No., Seed plant Name, and three columns for 1 hr, 3 hrs, 5 hrs, 10 hrs, and Control. Each time period column contains % germ, Shoot length (mm), and Root length (mm). Rows list Wheat, Jowar, Cowpea, Gram, Safflower, and Soyabea n.

References :

- 1. Ahmed I. Mehmood Z., Mohammad F. (1998). Screening of some medicinal plants for their antimicrobial properties.
2. Abdul- Baki A.A, and Anderson J.D. (1975) Vigour dermination in soyabea n seed by multiple criteria crop sci. B 630-633



3. Ali, A.A., K.E. Ali, A. Fadlalla and K.E. Khalid (2008) The effects of GA oral treatment on the metabolic profile of chronic renal failure patients under regular haemodialysis in Central Sudan. *Natural Product Research*, **22**(1): 12–21.
4. Ali, B.H. and M.S. Al Moundhri (2006) Agents ameliorating or augmenting the nephrotoxicity of cisplatin and other platinum compounds: a review of some recent research. *Food and Chemical Toxicology*, **44**(8): 1173–1183.
5. Babu, R., and Murugan, K. (1998). Interactive effect of neem seed kerna and neem gum extract on the control of *Culex quinquefasciatus* Say, *Neem Newsletter* 15(2): 9-11.
6. Badar, K.V. (2011) Studies on antimicrobial properties of gum of some plants. Ph.D. Thesis, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
7. Bliss, D.Z., T.P. Stein, C.R. Schleifer and R.G. Settle (1996) Supplementation with GA fiber increases fecal nitrogen excretion and lowers serum urea nitrogen concentration in chronic renal failure patients consuming a low-protein diet. *The American Journal of Clinical Nutrition*, **63**(3): 392–398.
8. Evans, A.J., R.L. Hood, D.G. Oakenfull and G.S. Sidhu (1992) Relationship between structure and function of dietary fibre: a comparative study of the effects of three galactomannans on cholesterol metabolism in the rat. *British Journal of Nutrition*, **68**(1): 217-229.
9. Glover, D.A., K. Ushida, A.O. Phillips and S.G. Riley (2009) Acacia(sen) SUPERGUMTM (Gum arabic): An evaluation of potential health benefits in human subjects. *Food Hydrocolloids*, **23**(8): 2410–2415.
10. Hinson, J.A., A.B. Reid, S.S. McCullough and L.P. James (2004) Acetaminophen-induced hepatotoxicity: role of metabolic activation, reactive oxygen/nitrogen species, and mitochondrial permeability transition. *Drug Metabolism Reviews*, **36**(3-4): 805–822.
11. Malcolm, D.W. (1936) Report on Gum and Gum arabic (Dar-es-salaam Gov. Printer). **3**(4): 188-191.
12. Matsumoto, N., S. Riley, D. Fraser, S. Al-Assaf, E. Ishimura, T. Wolever, G.O. Phillips and A.O. Phillips (2006) Butyrate modulates TGF-beta1 generation and function: potential renal benefit for *Acacia* (sen) SUPERGUM (G.A.)? *Kidney International*, **69**(2): 257–265.
13. Randall, R. C., G.O. Phillips and P.A. Williams (1988) The role of the proteinaceous component on the emulsifying properties of gum arabic. *Food Hydrocolloids*, **2**(2): 131–140.
14. Rehman, K.U., M.A. Wingertzahn, S. Teichberg, R.G. Harper and R.A. Wapnir (2003) Gum arabic. (GA) modifies paracellular water and electrolyte transport in the small intestine. *Digestive Diseases and Sciences*, **48**(4): 755–760.
15. Tiss, A., F. Carrière and R. Verger (2001) Effects of gum arabic on lipase interfacial binding and activity. *Analytical Biochemistry*, **294**(1): 36–43.
16. Trommer, H. and R.H. Neubert (2005) The examination of polysaccharides as potential antioxidative compounds for topical administration using a lipid model system. *International Journal of Pharmaceutics*, **298**(1): 153–163.
17. Wapnir, R.A., B. Sherry, C.N. Codipilly, L.O. Goodwin and I. Vancurova (2008) Modulation of rat intestinal nuclear factor NF-kappaB by gum arabic. Rat small intestine by gum arabic. *Digestive Diseases and Sciences*, **53**(1): 80–87.



-
18. Williams, P.A., G.O. Phillips and A.M. Stephen (1990) Spectroscopic and molecular comparisons of three fractions from Acacia senegal gum. *Food Hydrocolloids*, 4(4): 305-311.
 19. Zalucki M.P. and Malcollum S.B. (1999). Plants latex and first instar monarch larval growth and survival on three North Americal milkweed species. *Journal and chemical ecology* volume 25. Number 8, August 1999, PP-1827-1842 (16).

